

COHOMOLOGICAL PARTITION FUNCTIONS FOR A CLASS OF BOSONIC THEORIES

Antti J. Niemi* # and O. Tirkkonen

*Research Institute for Theoretical Physics, University of Helsinki
Siltavuorenpenger 20 C, SF-00170 Helsinki, Finland*

We argue that for a general class of nontrivial bosonic theories the path integral can be related to an equivariant generalization of conventional characteristic classes.

* E-mail: ANIEMI@PHCU.HELSENKI.FI

Permanent address after 1.7.1992: Institute for Theoretical Physics, Uppsala University, Uppsala, Sweden

During the last twenty years, characteristic classes [1] have been involved in several major developments both in physics and mathematics. The most familiar example of characteristic classes is the four dimensional $F\tilde{F}$ and its generalizations to other even dimensions, together with their odd-dimensional descendants such as the Chern-Simons action. Another widely encountered example is the gravitational $R\tilde{R}$ and its relatives. These characteristic classes are relevant to a large number of phenomena such as anomalies, fermion number fractionization, index theorems, CP-violation, fractional spin and statistics, high-temperature superconductivity and so on. More recently, the quantization of these characteristic classes has revealed several important results and provides us with new insight to problems from knot theory to quantum gravity [2].

In this Letter we wish to draw attention to a natural generalization of characteristic classes. This generalization has been discussed quite extensively in the contemporary mathematics literature, but until now it has not emerged in the physics literature. This generalization arises when ordinary cohomology is extended to equivariant cohomology [3-5]. In addition to the exterior derivative, equivariant cohomology also includes the action of a preferred vector field. In applications to physics such a preferred vector field is naturally present. It is the vector field that determines the hamiltonian flow. As a consequence equivariant cohomology provides a natural framework for extending conventional cohomology and its characteristic classes in a dynamically nontrivial fashion: Indeed, equivariant characteristic classes [4-5] share all the beauty, simplicity and power of their conventional counterparts, but have the additional feature that they account for a nontrivial hamiltonian action. Consequently we expect, that equivariant characteristic classes will become relevant to a number of important problems in physics.

Conventional characteristic classes are usually associated to the quantization of fermions: Anomalies are an important ingredient of the path integral measure in fermionic quantum field theories, and the Atiyah-Singer index theorem can also be derived from a supersymmetric path integral. In the present Letter we shall argue that equivariant characteristic classes can be associated to the quantization of *bosons*. Indeed, we shall identify a general class of bosonic hamiltonians such that the corresponding path integral evaluates to an equivariant characteristic class. The condition that we impose is quite general: Locally in a neighborhood, our condition is satisfied by a *generic* hamiltonian. Thus we expect that equivariant characteristic classes are an important ingredient also in the general quantization procedure, and corrections to our result seem to be related to the global aspects of the theory.

We shall first shortly explain aspects of equivariant cohomology. A more extensive discussion can be found *e.g.* in [5], and some applications to physics can be found *e.g.* in [6,7]. We shall begin from ordinary cohomology:

In ordinary cohomology we are interested in the action of a nilpotent exterior derivative d in the space of exterior forms which are associated to some manifold. In this space, d determines a linear mapping which sends an n -form onto an $n + 1$ -form. The n -forms which are annihilated by d are closed if they can not be represented as an exterior derivative of an $n - 1$ -form. Otherwise, they are exact. The space of closed

n -forms modulo exact n -forms determines the n th cohomology class of the manifold. Conventional characteristic classes are examples of nontrivial cohomology classes. In the physics literature, the most familiar is the Chern character of a gauge theory,

$$Ch(F) = Tr\{e^{\frac{1}{4\pi}F}\} \quad (1)$$

where F is the Yang-Mills curvature two-form in some (properly normalized) representation of the gauge group. Its gravitational analog is the $\hat{A}$ -genus

$$\hat{A}(R) = \prod_{i=1}^{\frac{n}{2}} \frac{\frac{1}{2}x_i}{sinh(\frac{1}{2}x_i)} \quad (2)$$

where the x_i are the skew-eigenvalues of the curvature two-form matrix R_{ab} . The expansion of these characteristic classes in their arguments yields closed forms, and the integrals of these closed forms are topological invariants of the manifold.

Equivariant cohomology is an extension of conventional cohomology. In addition of the exterior derivative, now there is also an interior multiplication (contraction) along a preferred vector field. With $\mathcal{X}^a$ components of the vector field, we denote the interior multiplication by

$$i_{\mathcal{X}} = \mathcal{X}^a i_a \quad (3)$$

where i_a is interior multiplication dual to some basis of one-forms $d\phi^a$ such that $i_a d\phi^b = \delta_a^b$. Interior multiplication maps an n -form onto an $n-1$ -form, and is nilpotent: $i_{\mathcal{X}}^2 = 0$. Combining exterior derivative with interior multiplication we then obtain the equivariant exterior derivative

$$d_{\mathcal{X}} = d + i_{\mathcal{X}} \quad (4)$$

Since d maps an n -form onto an $n+1$ -form and $i_{\mathcal{X}}$ maps an n -form onto an $n-1$ -form, $d_{\mathcal{X}}$ does not preserve the form degree but maps an even form onto an odd form, and vice versa. If we identify even forms with bosons and odd forms with fermions, we can then view $d_{\mathcal{X}}$ as a supersymmetry operator. The corresponding supersymmetry algebra closes to the Lie derivative along the vector field $\mathcal{X}^a$,

$$d_{\mathcal{X}}^2 = di_{\mathcal{X}} + i_{\mathcal{X}}d = \mathcal{L}_{\mathcal{X}} \quad (5)$$

and it has been established [7] that the supersymmetry algebra in a *generic* supersymmetric theory is of the form (5) in the (super)loop space of the theory, provided one introduces an appropriate auxiliary field formalism.

The equivariant exterior derivative (4) determines an equivariant generalization of cohomology on the manifold. One can show that this equivariant cohomology can be identified with conventional cohomology on the subspace of exterior forms which are annihilated by the Lie derivative (5). Indeed, in this subspace the equivariant exterior derivative (4) becomes nilpotent, and consequently it determines a conventional exterior derivative in this subspace. Representatives of nontrivial cohomology classes can then be constructed using the pertinent conventional characteristic classes. However, since the equivariant exterior derivative involves interior multiplication along the

vector field $\mathcal{X}^a$, these characteristic classes necessarily contain this vector field in a definite, but nontrivial fashion. Furthermore, since equivariant cohomology coincides with ordinary cohomology in the subspace where (5) vanishes, it is obvious that in spite of their nontrivial dynamical content, equivariant characteristic classes share the geometrical properties of their conventional counterparts and can be similarly analysed using powerful geometrical techniques [5].

The purpose of this Letter is to draw attention to equivariant characteristic classes as potentially important quantities in physics. In order to exemplify this, we shall now proceed to a path integral derivation of the equivariant generalizations of (1) and (2). We shall find that *for a very general class of bosonic hamiltonians, the path integral evaluates to a combination of the equivariant Chern character and the equivariant $\hat{A}$ -genus, i.e.* an equivariant generalization of the Atiyah-Singer index density for a Dirac operator. For this, we consider a phase space with a generic coordinate system ϕ^a . Here the index a may be either finite or infinite, and in a quantum field theory application it denotes both internal degrees of freedom and space coordinates. The symplectic two-form on this phase space is

$$\omega = \frac{1}{2} \omega_{ab} d\phi^a \wedge d\phi^b \quad (6)$$

and the inverse matrix ω^{ab} determines Poisson brackets by

$$\{A(\phi), B(\phi)\} = \omega^{ab} \partial_a A \partial_b B \quad (7)$$

Since $d\omega = 0$, we can locally represent it as an exterior derivative of the symplectic one-form $\theta = \theta_a d\phi^a$,

$$\omega = d\theta \quad (8)$$

We are interested in the equivariant cohomology determined by the canonical action of a given hamiltonian H . For this, we introduce the corresponding hamiltonian vector field $\mathcal{X}_H$ which is defined by

$$\mathcal{X}_H^a = \omega^{ab} \partial_b H \quad (9)$$

We can then introduce the corresponding equivariant exterior derivative

$$d_H = d + i_H \quad (10)$$

and the Lie-derivative along (9)

$$\mathcal{L}_H = di_H + i_H d \quad (11)$$

In the following we shall *assume* that the phase space admits a metric tensor g_{ab} which is Lie-derived by $\mathcal{X}_H$,

$$\mathcal{L}_H g = 0 \quad (12)$$

Locally we can always find this metric tensor in a small enough phase space neighborhood (and outside of the critical points of H) but there might be global difficulties

associated in defining g_{ab} . Consequently (12) determines a restriction on the class of hamiltonians that we shall consider. However, since (12) can be satisfied by a large number of interesting hamiltonians defined *e.g.* on phase spaces with a Kähler structure, we do not view (12) as an overly strong restriction.

We are interested in evaluating the phase space path integral

$$Z = \int [d\phi^a] \sqrt{\text{Det}(\omega_{ab})} \exp\left\{i \int_0^T \theta_a \dot{\phi}^a - H\right\} \quad (13)$$

with periodic boundary conditions. For this, we first introduce anticommuting variables c^a and write (13) as

$$Z = \int [d\phi^a][dc^a] \exp\left\{i \int_0^T \theta_a \dot{\phi}^a - H + \frac{1}{2} c^a \omega_{ab} c^b\right\} \quad (14)$$

In [6] we have investigated path integrals of the form (14). There, we interpret (14) as a loop space integral, with the loop space parametrized by the time evolution $\phi^a \rightarrow \phi^a(t)$. The bosonic part of the action then determines a loop space hamiltonian vector field,

$$\mathcal{X}_S^a = \dot{\phi}^a - \omega^{ab} \partial_b H \quad (15)$$

and we can identify $c^a(t)$ as the basis of one-forms on this loop space. Defining the loop space exterior derivative (in the following time integrals are implicit)

$$d = c^a \partial_a \quad (16)$$

we then obtain the loop space version of the equivariant exterior derivative,

$$d_S = c^a \partial_a + \mathcal{X}_S^a i_a \quad (17)$$

Here i_a is a basis for loop space interior multiplication, defined by $i_a c^b = \delta_a^b$.

In analogy with (10), (11) we conclude that the equivariant exterior derivative (17) determines a loop space supersymmetry transformation. The path integral (14) is (formally) invariant under this supersymmetry transformation. From this we conclude [6] that if ψ is a loop space one-form such that

$$\mathcal{L}_S \psi = (di_S + i_S d) \psi = 0 \quad (18)$$

then the path integral

$$Z_\psi = \int [d\phi^a][dc^a] \exp\left\{i \int_0^T \theta_a \dot{\phi}^a - H + \frac{1}{2} c^a \omega_{ab} c^b + d_S \psi\right\} \quad (19)$$

is (formally) independent of ψ , and coincides with the original path integral (14). In particular, if we select ψ properly we obtain the path integral generalization [6] of the

Duistermaat-Heckman integration formula [5,8], which states that the path integral (19) localizes to the critical points of the bosonic action. Alternative choices of ψ then yield alternative integration formulas corresponding to localizations to different field configurations. If more than one integration formula is valid for a given theory, their answers must coincide. However, for some theories it may happen that some integration formulas are not valid. Then, different integration formulas can yield different results and we must select the one which gives the correct result, if it exists.

In the present Letter we shall employ the condition (12) to introduce a class of functionals ψ which is *different* from the one used in [6]. We then obtain a localization of the path integral (19) to field configurations which are *different* from the configurations that are relevant in the Duistermaat-Heckman integration formula: In particular, we find that our localization provides an integration formula for the path integral (14) in terms of equivariant characteristic classes.

If the condition (12) is satisfied, the following one-parameter family of one-forms is in the subspace (18),

$$\psi_\lambda = \frac{\lambda}{2} g_{ab} \dot{\phi}^a c^b \quad (20)$$

The corresponding path integral

$$Z_\lambda = \int [d\phi^a][dc^a] \exp\left\{i \int_0^T \theta_a \dot{\phi}^a - H + \frac{1}{2} c^a \omega_{ab} c^b + \frac{\lambda}{2} d_S(g_{ab} \dot{\phi}^a c^b)\right\} \quad (21)$$

is then (formally) independent of λ , and in the $\lambda \rightarrow 0$ limit it reduces to the original path integral (14). In the following we shall be interested in the $\lambda \rightarrow \infty$ limit of (21):

Explicitly, the action in (21) reads

$$S = \frac{\lambda}{2} g_{ab} \dot{\phi}^a \dot{\phi}^b + (\theta_a - \frac{\lambda}{2} g_{ab} \mathcal{X}_H^b) \dot{\phi}^a - H + \frac{\lambda}{2} c^a (g_{ab} \partial_t + \dot{\phi}^c g_{bd} \Gamma_{ac}^d) c^b + \frac{1}{2} c^a \omega_{ab} c^b \quad (22)$$

Here Γ_{ac}^d is the Christoffel symbol for the metric tensor g_{ab} . Since the path integral (21) is (formally) independent of λ , we can evaluate it in the $\lambda \rightarrow \infty$ limit. For this, we introduce

$$\phi^a(t) = \phi_0^a + \phi_t^a \quad (23.a)$$

$$c^a(t) = c_0^a + c_t^a \quad (23.b)$$

with ϕ_0^a, c_0^a the constant modes of $\phi^a(t)$ and $c^a(t)$, and ϕ_t^a, c_t^a the t -dependent modes. We then define the path integral measure as

$$[d\phi^a][dc^a] = d\phi_0^a dc_0^a \prod_t d\phi_t^a dc_t^a \quad (24)$$

and introduce the change of variables

$$\phi_t^a \rightarrow \frac{1}{\sqrt{\lambda}} \phi_t^a \quad (25.a)$$

$$c_t^a \rightarrow \frac{1}{\sqrt{\lambda}} c_t^a \quad (25.b)$$

Formally, the Jacobian for this change of variables is trivial. In the $\lambda \rightarrow \infty$ limit we can then evaluate the path integral over the non-constant modes, which yields

$$Z = \int \sqrt{g} d\phi_0^a d c_0^a \frac{e^{-iTH + i\frac{T}{2} c_0^a \omega_{ab} c_0^b}}{\sqrt{\text{Det}[g_{ab} \partial_t - \frac{1}{2}(\Omega_{ab} + R_{abcd} c_0^c c_0^d)]}} \quad (26)$$

Here we have defined

$$\Omega_{ab} = \partial_b(g_{ac} \mathcal{X}_H^c) - \partial_a(g_{bc} \mathcal{X}_H^c) \quad (27)$$

and $R_{abcd} c_0^c c_0^d$ is the curvature two-form. We evaluate the determinant *e.g.* by the ζ -function method. This gives

$$Z = \int \sqrt{g} d\phi_0^a d c_0^a e^{-iTH + i\frac{T}{2} c_0^a \omega_{ab} c_0^b} \sqrt{\text{Det} \left[\frac{\frac{T}{2}(\Omega_{ab} + R_{ab})}{\sinh[\frac{T}{2}(\Omega_{ab} + R_{ab})]} \right]} \quad (28)$$

Notice that if we set $H = 0$, the exponential in (28) yields the Chern character for the two-form $c_0^a \omega_{ab} c_0^b$, while the second term in (28) gives the $\hat{A}$ -genus for the curvature two-form $R_{abcd} c_0^c c_0^d$. This coincides with the result one expects from a path integral evaluation of the Atiyah-Singer index theorem for a Dirac operator in the background of a U(1) gauge field and a gravitational field. Indeed, for $H = 0$ the action in (28) reduces to

$$S \rightarrow \frac{\lambda}{2} g_{ab} \dot{\phi}^a \dot{\phi}^b + \theta_a \dot{\phi}^a + \frac{\lambda}{2} c^a (g_{ab} \partial_t + \dot{\phi}^c \Gamma_{acb}) c^b + c^a \omega_{ab} c^b \quad (29)$$

which is identical to the action used in [7] to evaluate the Atiyah-Singer index theorem, provided we identify θ_a and ω_{ab} with the gauge field A_μ and its field strength tensor $F_{\mu\nu}$.

We shall now argue, that with nontrivial H the two terms in (28) are equivariant generalizations of the Chern character (1) *resp.* $\hat{A}$ -genus (2) in the equivariant cohomology determined by the phase space equivariant exterior derivative (10). Indeed, since

$$d_H(H - \frac{1}{2} c^a \omega_{ab} c^b) = 0 \quad (30)$$

we can immediately identify the exponential term in (28) as an equivariant generalization of the Chern-character (1). Furthermore, if we introduce the covariant generalization of the equivariant exterior derivative

$$D_H = d + \Gamma + i_H \quad (31)$$

where Γ is the Christoffel symbol one-form, we find that

$$D_H(\Omega_{ab} + R_{ab}) = 0 \quad (32)$$

From this we can conclude that the second term in (28) is also equivariantly closed,

$$(d + i_H) \sqrt{\text{Det} \left[\frac{\frac{T}{2}(\Omega_{ab} + R_{ab})}{\sinh[\frac{T}{2}(\Omega_{ab} + R_{ab})]} \right]} = 0 \quad (33)$$

Hence it determines an equivariant generalization of the $\hat{A}$ -genus. As a consequence we have verified that formally *the bosonic phase space path integral (14) evaluates to a combination of the equivariant Chern character and the equivariant $\hat{A}$ -genus, i.e. an equivariant generalization of the Atiyah-Singer index density for a Dirac operator.*

The equivariant $\hat{A}$ -genus in (28) is determined by expanding $\hat{A}$ at $\frac{T}{2}\Omega_{ab}$ in the "variable" $\frac{T}{2}R_{ab}$. Since the function

$$f(z) = \frac{\frac{z}{2}}{\sinh \frac{z}{2}} \quad (34)$$

has poles at $2ik\pi$, this suggests an interesting phase structure for the pertinent thermodynamical partition function at finite temperatures $T \rightarrow i\beta$.

We observe that our final result (28) is quite similar to the Lefschetz formulas by Atiyah, Bott, Singer [9] and Bismuth [10]. Indeed, if in addition of (12) we also assume that the hamiltonian Lie-derives the symplectic one-form,

$$\mathcal{L}_H \theta = 0 \quad (35)$$

we can use loop space equivariant cohomology to write the entire action (29) in the form

$$S = (d + i_{\dot{\phi}} + i_H) \left(\frac{\lambda}{2} g_{ab} \dot{\phi}^a c^b + \theta_a c^a \right) \quad (36)$$

Here we have defined

$$i_{\dot{\phi}} = \dot{\phi}^a i_a \quad (37)$$

and i_H is now interpreted as a loop space interior multiplication. This representation of the action admits the following interpretation: As before, we can identify

$$(d + i_{\dot{\phi}}) \left(\frac{\lambda}{2} g_{ab} \dot{\phi}^a c^b + \theta_a c^a \right) = \frac{\lambda}{2} g_{ab} \dot{\phi}^a \dot{\phi}^b + \theta_a \dot{\phi}^a + \frac{\lambda}{2} c^a (g_{ab} \partial_t + \dot{\phi}^c \Gamma_{acb}) c^b + c^a \omega_{ab} c^b \quad (38)$$

with a Dirac operator in the background of a U(1) gauge field θ_a and a gravitational field. If we denote the chiral components of this Dirac operator by D_+ and D_- , the remaining terms in (36) then coincide with the terms, that we expect to obtain in a supersymmetric path integral representation of the Lefschetz number

$$L(e^{iTH}) = \text{Tr} \{ e^{iTH} (e^{-iTD_+ D_-} - e^{-iTD_- D_+}) \} \quad (39)$$

for the hamiltonian H . As a consequence we conclude that in the case of hamiltonians for which the conditions (12), (35) are satisfied, our result (28) reproduces the result by Atiyah, Bott, Singer [9] and Bismuth [10].

From the relation between (14) and (39) we can draw the following conceptually interesting interpretation of our result: We have found, that a bosonic path integral (14) can be related to the properties of a (functional) Dirac operator defined in the canonical phase space of the bosonic theory. In particular, the path integral (essentially) coincides with the Lefschetz number of the bosonic hamiltonian evaluated for this Dirac operator. (Notice that in field theory applications this Dirac operator is defined in an infinite dimensional space.)

In conclusion, we have investigated a general family of bosonic path integrals. We have found that these path integrals can be evaluated in a closed form, and the result coincides with the equivariant generalization of the Atiyah-Singer index density for a Dirac operator. The restriction (12) that we have introduced is relatively mild - locally, it can be satisfied by an arbitrary hamiltonian. This indicates, that the corrections to our result should in some sense reflect global properties of the theory. Our computations have also been formal in that we have overlooked *e.g.* difficulties associated with the definition of the path integral measure: There might be anomalies associated with our argument that the path integral is λ independent. However, since we have also obtained a relation between our result and the (infinitesimal) Lefschetz formula by Atiyah, Bott, Singer and Bismuth, we expect that in a number of interesting cases our arguments can be made rigorous. Finally, we conclude that our result clearly indicates that equivariant characteristic classes can be highly relevant to a number of important problems in theoretical physics.

A.N. thanks J.M. Bismuth and A. Reyman for discussions. Both authors thank the Theory Division at CERN for hospitality during this work. A.N. also thanks Ecole Polytechnique and Paris VI for hospitality during this work.

References

1. For a review, see T. Eguchi, P.B. Gilkey and A.J. Hanson, Phys. Repts. **66** (1980) 213
2. E. Witten, Comm. Math. Phys. **117** (1988) 353; For a review, see D. Birmingham, M. Blau, M. Rakowski and G. Thompson, Phys. Repts. **209** (1991) 129
3. E. Witten, J. Diff. Geom. **17** (1982) 661
4. N. Berline and M. Vergne, Duke Math. Journ. **50** (1983) 539
5. For a review, see N. Berline, E. Getzler and M. Vergne, *Heat Kernels and Dirac Operators* (Springer-Verlag, Berlin, 1991)
6. M. Blau, E. Keski-Vakkuri and A.J. Niemi, Phys. Lett. **B246** (1990) 92; A.J. Niemi and P. Pasanen, Phys. Lett. **B253** (1991) 349; E. Keski-Vakkuri, A.J. Niemi, G. Semenoff and O. Tirkkonen, Phys. Rev. **D44** (1991) 3899
7. A. Hietamäki, A.Yu. Morozov, A.J. Niemi and K. Palo, Phys. Lett. **B263** (1991) 417; A. Yu. Morozov, A.J. Niemi and K. Palo, Physics Letters **B271** (1991) 365; Nucl. Phys. **B** (to appear)
8. J.J. Duistermaat and G.J. Heckman, Inv. Math. **72** (1983) 153; N. Berline and M. Vergne, Amer. J. Math, **107** (1985) 1159
9. M. Atiyah and R. Bott, Ann. Math. **86** (1967) 374; *ibid.* **88**, (1968) 451; M. Atiyah and I. Singer, Ann. Math. **87** (1968) 546; *ibid.* **93** (1971) 119
10. J.-M. Bismut, Comm. Math. Phys. **98** (1985) 213; *ibid.* **103** (1986) 127; J. Funct. Anal. **62** (1985) 435